

Just One Bite® EX Bait Station

Farnam Companies, Inc.

Description:

Kills House Mice. This bait station is not tamper resistant. Use indoors Only in areas inaccessible to children and pets.

*House Mice usually consume a **lethal dose** in a single feeding, but it may take two or more days from time of bait consumption for them to die.

Due to **Stop Feed Action**, house mice cease feeding after consuming a toxic dose.

This product is effective against anticoagulant-resistant house mice.

Effective Indoor Mouse Control

ACTIVE INGREDIENT:

Bromethalin (CAS No. 63333-35-7)
..... 0.01%

OTHER
INGREDIENTS:.....99.
99%

TOTAL:.....
.....100.00%

Contains Bitrex

KEEP OUT OF REACH OF CHILDREN

MAY BE HARMFUL IF EATEN BY CHILDREN OR PETS

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

CAUTION: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation.

Keep away from children, domestic animals and pets. Do not get in eyes, on skin, or on clothing. All handlers (including applicators) must wear long-sleeved shirt and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

User Safety Requirements

As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco, or using the toilet, and change into clean clothing.

FIRST AID

Have label with you when obtaining treatment advice

IF SWALLOWED: Call a poison control center or doctor or 1-800-234-2269 immediately for

treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting

unless told to do so by the poison control center or doctor. Do not give anything by mouth to an

unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin

immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for

treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20

minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice.

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian at once.

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the nerve poison Bromethalin. This is not an anticoagulant type rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to mammals and birds. Dogs and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water.

Important:

This is not a tamper-resistant bait station. Do not use this product outdoors or in areas accessible to children, pets, domestic animals, and nontarget wildlife. To help prevent accidents:

1. Store product in use in a location out of reach of children and pets.
2. Dispose of product container and unused, spoiled and unconsumed bait as specified on this label.

USE SITES: For indoor use only. This product may only be used to control house mice in homes and residential buildings.

USE RESTRICTIONS: Do not use this product in sewers or any outdoor area. Do not place this ready-to-use bait station in any area where there is a possibility of contaminating food or surfaces that come into direct contact with food. Do not break station open and/or deliberately remove bait from it. Do not sell this product in individual containers holding more than 1 pound of bait.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where house mice or their signs have been seen. Remove as much alternative food accessible to rodents as possible.

APPLICATION DIRECTIONS: The active ingredient in this product is different from anticoagulant rodenticides. Individual house mice cease feeding after consuming a toxic dose.

HOUSE MICE: Wearing gloves, place bait stations 8 to 12 feet apart in infested areas. Highest bait consumption is expected to occur on the first day or two after treatment. After several days, while wearing gloves, inspect bait placements and replace with new bait stations at sites where there is heavy feeding or where there is continued evidence of house mice activity. Continue baiting until all signs of house mouse activity have ceased.

Wearing gloves, collect and properly dispose of all dead house mice and used bait stations. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.