

NDC 0010-3832-02

# Synanthic®

(oxfendazole)

## Bovine Dewormer Suspension

For Animal Use Only

NADA 140-854, Approved by FDA



Efficacy comparison:						
	<i>Haemonchus</i>		<i>Ostertagia</i>		<i>Cooperia</i>	
	FECRT*	Adult Control	FECRT*	Adult Control	FECRT*	Adult Control
<b>Synanthic® (oxfendazole)</b>	100%	98%	94%	90%	100%	99%
<b>Fenbendazole</b>	99%	100%	65%	73%	99%	99%

\*Fecal egg count reduction test

	Brown Stomach Worm			
	Adult	L <sub>4</sub>	Inhibited L <sub>4</sub>	Tapeworm
<b>Synanthic® 4.5 mg/kg (oxfendazole)</b>	●	●	●	●
<b>Fenbendazole (5 mg/kg)</b>	●			
<b>Fenbendazole (10mg/kg)</b>	●	●	●	●

### SYNANTHIC® RESIDUE WARNING:

Cattle must not be slaughtered until seven days after treatment. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. There are no contraindication for the use of Synanthic bovine dewormer suspension in beef cattle.